



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>A61K 31/00</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/59487</b> <b>(43) International Publication Date:</b> 12 October 2000 (12.10.00)
<b>(21) International Application Number:</b> PCT/US00/06446 <b>(22) International Filing Date:</b> 5 April 2000 (05.04.00)  <b>(30) Priority Data:</b> 60/128,141 7 April 1999 (07.04.99) US 09/534,325 24 March 2000 (24.03.00) US  <b>(71)(72) Applicant and Inventor:</b> STOLL, Andrew, L. [US/US]; 35 Old Winter Street, Lincoln, MA 01773 (US).  <b>(74) Agents:</b> SANZO, Michael, A. et al.; Pillsbury Madison & Sutro, LLP, 1100 New York Avenue, N.W., Washington, DC 20005 (US).		<b>(81) Designated States:</b> AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> FLUPIRTINE IN THE TREATMENT OF FIBROMYALGIA AND RELATED CONDITIONS  <b>(57) Abstract</b>  The present invention is directed to a method for treating the symptoms associated with fibromyalgia and related conditions by administering flupirtine.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

## Flupirtine in the Treatment of Fibromyalgia and Related Conditions

### Field of the Invention

- 5 The present invention is directed to medical treatments for fibromyalgia and related conditions. Specifically, the invention is directed to the administration of the drug flupirtine as a means for alleviating the symptoms associated with these disorders.

### Background of the Invention

- 10 Fibromyalgia is a chronic condition characterized by pain in muscles, fascia and joints. Other symptoms typically include sleep disturbances, chronic fatigue and major depression. The etiology and pathophysiology of fibromyalgia are unknown, but it is clear that the central nervous system is involved. Patients may obtain a degree of relief from analgesic drugs, antidepressants and adjunctive treatments such as moderate exercise, proper diet and stress reduction techniques.
- 15 Table 1 summarizes the studies that have been carried out in an effort to find an effective drug treatment.

**Table 1: Drug Treatments for Fibromyalgia**

Treatment	Drug Class*	Study Method**	Efficacy	Reference
fluoxetine	1	open	+	Johnson, <i>J. Fam. Pract.</i> , 44:128-30 (1997)
fluoxetine	1	open	+	Finestone, <i>et al.</i> , <i>J.A.M.A.</i> , 264:2869-70 (1990)
fluoxetine	1	open	+	Goldenberg, <i>et al.</i> , <i>Arthritis Rheumatol.</i> , 39:1852-9 (1996)
fluoxetine	1	db	±	Wolfe, <i>et al.</i> , <i>Scand. J. Rheumatol.</i> , 23:255-9 (1994)
fluoxetine	1	open	-	Cortet, <i>et al.</i> , <i>Rev. Rheum. Mal. Osteoartic</i> , 59:497-500 (1992)
citalopram	1	db	-	Norregaard, <i>et al.</i> , <i>Pain</i> ,

Treatment	Drug Class*	Study Method**	Efficacy	Reference
				61:445-9 (1995)
amitriptyline	1	db	+	Carette, <i>et al.</i> , <i>Arthritis Rheum.</i> , 37:32-40 (1994)
amitriptyline	1	db	+	Carette, <i>et al.</i> , <i>Arthritis Rheum.</i> , 29:655-9 (1986)
amitriptyline	1	db	+	Goldenberg, <i>et al.</i> , <i>Arthritis Rheumatol.</i> , 39:1852-9 (1996)
amitriptyline	1	db	+	Goldenberg, <i>et al.</i> , <i>Arthritis Rheum.</i> , 29:1371-7 (1986)
amitriptyline	1	db	+	Jaeschke, <i>et al.</i> , <i>J. Rheumatol.</i> , 18:447-51 (1991)
amitriptyline	1	db	+	Scudds, <i>et al.</i> , <i>J. Rheumatol.</i> , 16:98-103 (1989)
fluphenazine plus amitriptyline	7, 1	open	+	Connolly, <i>Del. Med. J.</i> , 53:189-91 (1981)
naproxen plus amitriptyline	1, 2	db	+	Goldenberg, <i>et al.</i> , <i>Arthritis Rheum.</i> , 29:1371-7 (1986)
ondansetron	9	db	+	Hrycaj, <i>et al.</i> , <i>J. Rheumatol.</i> , 23:1418-23 (1996)
tropisetron	9	open	+	Samborski, <i>et al.</i> , <i>Mater Med. Pol.</i> , 28:17-9 (1996)
ketanserin	9	open	+	Stratz, <i>et al.</i> , <i>Z. Rheumatol.</i> , 50:21-2 (1991)
lithium carbonate	6	open		Teasell, <i>Can. Med. Assoc. J.</i> , 144:122-3 (1991)
lithium carbonate	6	open		Tyber, <i>Can. Med. Assoc. J.</i> , 143:902-4 (1990)
fluoxetine plus cyclobenzaprine	1, 3	open	+	Cantini, <i>et al.</i> , <i>Minerva Med.</i> , 85:97-100 (1994)
cyclobenzaprine	3	db	+	Carette, <i>et al.</i> , <i>Arthritis Rheum.</i> , 37:32-40 (1994)
cyclobenzaprine	3	db	+	Bennett, <i>Arthritis Rheum.</i> ,

Treatment	Drug Class*	Study Method**	Efficacy	Reference
				31:1535-42 (1988)
cyclobenzaprine	3	db	+	Quimby, et al., <i>J. Rheumatol. Suppl.</i> , 19:140-3 (1989)
low-dose hs cyclobenzaprine	3	db	+	Santandrea, et al., <i>J. Int. Med. Res.</i> , 21:74-80 (1993)
cyclobenzaprine plus ibuprofen	3, 2	open	+	Fossaluzza, et al., <i>Int. J. Clin. Pharmacol. Res.</i> , 12:99-102 (1992)
carisoprodol plus acetaminophen plus caffeine	3, 12, 13	db	+	Vaeroy, et al., <i>Clin. Rheumatol.</i> , 8:245-50 (1989)
Zolpidem	4	db	±	Moldofsky, et al., <i>J. Rheumatol.</i> , 23:529-33 (1996)
ibuprofen plus alprazolam	4	db	±	Russell, et al., <i>Arthritis Rheum.</i> , 34:552-60 (1991)
ketamine	15	open	+	Oye, et al., <i>Tidsskr. Nor. Laegeforen</i> , 116:3130-1 (1996)
indomethacin	2	open	+	Kelly, <i>J. Am. Geriatr. Soc.</i> , 14:48-55 (1966)
naproxen	2	db	±	Goldenberg, et al., <i>Arthritis Rheum.</i> , 29:1371-7 (1986)
tenoxicam	2	db	+	Schorn, <i>S. Afr. Med. J.</i> , 69:301-3 (1986)
ibuproxam	2	open	+	Ferri, et al., <i>Minerva Med.</i> , 74:331-6 (1983)
tolmetin	2	db	+	Balme, et al., <i>Curr. Med. Res. Opin.</i> , 7:127-30 (1980)
tiaprofenic acid	2	db	+	Donald, et al., <i>J. Int. Med. Res.</i> , 8:382-7 (1980)
aspirin	2	db	+	Donald, et al., <i>J. Int. Med. Res.</i> , 8:382-7 (1980)
ibuprofen	2	db	+	Le Gallez, et al., <i>Curr. Med.</i>

Treatment	Drug Class*	Study Method**	Efficacy	Reference
				<i>Res. Opin.</i> , 10:663-7 (1988)
ibuprofen	2	db	-	Yunus, <i>et al.</i> , <i>J. Rheumatol.</i> , 16:527-32 (1989)
ibuprofen plus meptazinol	2, 9	db	+	Le Gallez, <i>et al.</i> , <i>Curr. Med. Res. Opin.</i> , 10:663-7 (1988)
phenylbutazone	2	db	+	Balme, <i>et al.</i> , <i>Curr. Med. Res. Opin.</i> , 7:127-30 (1980)
phenylbutazone plus prednisone	2, 8	open		Settel, <i>Curr. Ther. Res. Clin. Exp.</i> , 9:197-9 (1967)
topical capsaicin	10	open	+	Mathias, <i>et al.</i> , <i>Am. J. Phys. Med. Rehabil.</i> , 74:39-44 (1995)
local trigger point injections	11	open	±	Hong, <i>et al.</i> , <i>Arch. Phys. Med. Rehabil.</i> , 77:1161-6 (1996)
local trigger point injections	11	open		Jayson, <i>Bull. Hosp. Jt. Dis.</i> , 55:176-7 (1996)
antibiotics specific for Lyme disease	5	open	-	Dinerman, <i>et al.</i> , <i>Ann. Intern. Med.</i> , 117:281-5 (1992)
prednisone	8	db	-	Clark, <i>et al.</i> , <i>J. Rheumatol.</i> , 12:980-3 (1985)
chlormezanone	16	db	-	Patrick, <i>et al.</i> , <i>Br. J. Rheumatol.</i> , 32:55-8 (1993)
proglumetacin	16	open	+	Gusso, <i>et al.</i> , <i>Minerva Med.</i> , 76:567-73 (1985)
acetaminophen	12	db	-	Hrycaj, <i>et al.</i> , <i>J. Rheumatol.</i> , 23:1418-23 (1996)
s-adenosyl-methionine	16	db	+	Jacobsen, <i>et al.</i> , <i>Scand. J. Rheumatol.</i> , 20:294-302 (1991)
zopiclone	16	db	±	Drewes, <i>et al.</i> , <i>Scand. J. Rheumatol.</i> , 20:288-93 (1991)
zopiclone	16	db	-	Gronblad, <i>et al.</i> , <i>Clin. Rheumatol.</i> , 12:186-91 (1993)

Treatment	Drug Class*	Study Method**	Efficacy	Reference
intravenous lignocaine	14	open	+	Bennett, <i>et al.</i> , <i>Int. J. Clin. Pharmacol. Res.</i> , 15:115-9 (1995)
5-OH-L-tryptophan	16	open	+	Puttini, <i>et al.</i> , <i>J. Int. Med. Res.</i> , 20:182-9 (1992)
malic acid + magnesium	16, 16	db	+	Russell, <i>et al.</i> , <i>J. Rheumatol.</i> , 22:953-8 (1995)
tamoxifen	16	open		Simonson, <i>Lakartidningen</i> , 93:340 (1996)
botulinum toxin	16	open	-	Paulson, <i>et al.</i> , <i>Mov. Disord.</i> , 11:459 (1996)

\*Drug class key: 1 = antidepressant; 2 = NSAIDs; 3 = muscle relaxants (nonbenzodiazepine); 4 = benzodiazepine; 5 = antibiotics; 6 = mood stabilizer; 7 = antipsychotic agent; 8 = corticosteroid; 9 = serotonin (2 or 3) receptor antagonist; 10 = topical agents; 11 = trigger point injection; 12 = acetaminophen; 13 = caffeine; 14 = intravenous anesthetics; 15 = NMDA (glutamate) receptor ligand; 16 = miscellaneous agents.

\*\*Study method: db = double-blind

Despite the efforts that have been made, there is still no treatment that is effective in the majority of patients with fibromyalgia. Thus, there is a clear need for new therapies designed to alleviate the suffering of patients with this, and closely related, conditions.

## 15 Summary of the Invention

Flupirtine is an analgesic that has been used in Europe to treat the pain associated with surgery, cancer, trauma, and liver disease. It acts via the central nervous system through non-opiate pain pathways, possibly involving the thalamus or spinal pain pathways. In some, but not all, studies flupirtine has been found to be as effective as opiates in relieving pain. Moreover, flupirtine offers a clear advantage over opiates in that it is not addictive and there have been no reports of abuse. The present invention is based upon the unexpected discovery that flupirtine is capable of completely eliminating the chronic and severe body pain present in some patients with

fibromyalgia. Relief is possible even when high doses of opiates, non-steroidal anti-inflammatory drugs and placquenil fail to alleviate pain.

In its first aspect, the invention is directed to a method of treating a patient with fibromyalgia by administering a pharmaceutical composition containing flupirtine. The dosage of flupirtine should be sufficient to reduce or eliminate one or more of the symptoms that have been associated with fibromyalgia. These symptoms include diffuse musculoskeletal pain, fatigue, ocular and vestibular disturbances, headaches, paresthesias, esophageal dysmotility, sleep disturbances, and severe depression. The flupirtine may be administered in any dosage form including tablets, capsules, gels, or topical preparations and should be given at a daily dosage of between 50 and 900 mg, calculated on the basis of the free base form of the drug. In general, dosage should not exceed 600 mg per day and the optimal range should typically be between 100 and 400 mg per day. The flupirtine may be administered in either a single or, preferably, multidosage daily regimen.

In addition, the present invention is directed to methods of treating a patient for diseases or conditions that are related to fibromyalgia by administering flupirtine. Again, the dosage should be between 50 and 900 mg per day, calculated on the basis of the free base form of the drug, and should preferably be between 100 and 400 mg per day. The flupirtine may be given in any dosage form and administration may be accomplished by any route. Specific diseases that are defined as being related to fibromyalgia include fibrositis; chronic fatigue syndrome; myofascial pain syndrome, soft tissue rheumatism; idiopathic muscle pain syndrome; chronic widespread musculoskeletal pain; major depression associated with musculoskeletal pain; somatization disorder; somatoform pain disorder; Lyme disease with concurrent fibromyalgia; rheumatoid arthritis with concurrent fibromyalgia; generalized tendomyopathy; eosinophilia-myalgia syndrome; and stiff-man syndrome.

#### **Detailed Description of the Invention**

Fibromyalgia is one of the most common conditions seen by rheumatologists. Strict criteria for the diagnosis and classification of fibromyalgia have been established by, *inter alia*, the American College of Rheumatology. The main clinical characteristic is generalized pain and



widespread tenderness on palpation in specific areas of the musculoskeletal system. Stiffness, chronic fatigue and a specific sleep disturbance are also commonly present along with a number of other symptoms. The condition may be either idiopathic (primary fibromyalgia) or it may be associated with other underlying conditions such as ankylosing spondylitis, trauma or surgery (secondary fibromyalgia).

Flupirtine, a triaminopyridine compound with antinociceptive effects, has been marketed in Europe for several years by Asta Medica under the brand name Katadolon™. Although it has been used as an analgesic for cancer pain, post-operative pain, dental pain, degenerative rheumatic arthrosis, and inflammatory rheumatoid arthritis, it has not previously been known to be useful as a treatment for fibromyalgia. The present invention is based upon the discovery that flupirtine is unexpectedly effective in alleviating symptoms of fibromyalgia.

#### *Chemical Form of Flupirtine*

The present invention is not limited to any particular chemical form of flupirtine and the drug may be given to patients either as a free base or as a pharmaceutically acceptable acid addition salt. In the latter case, the hydrochloride salt is generally preferred but other salts derived from organic or inorganic acids may be also used. Examples of such acids include, without limitation, hydrobromic acid, phosphoric acid, sulfuric acid, methane sulfonic acid, phosphorous acid, nitric acid, perchloric acid, acetic acid, tartaric acid, lactic acid, succinic acid, citric acid, malic acid, maleic acid, aconitic acid, salicylic acid, thalic acid, embonic acid, enanthic acid, and the like. The preparation of flupirtine, 2-amino-3-carbethoxyamino-6-(4-fluorobenzylamino)-pyridine, and its physiologically acceptable salts is described in German patents 1,795,858 and 3,133,519.

#### *Dosage*

The total daily dosage of flupirtine administered to a patient should be at least the amount required to reduce or eliminate one or more of the symptoms associated with fibromyalgia, preferably musculoskeletal pain. The typical daily dosage will be between 100 and 300 mg and, in general, the daily dosage should not exceed 600 mg. Higher doses may be tolerated by some patients and daily dosages of 1,000 mg or more may be considered in refractory cases or in

patients receiving concomitant drug treatment with agents that may lower the serum concentration and half-life of flupirtine (e.g., cytochrome P450 inducing compounds such as carbamazepine, phenytoin, phenobarbital and rifampin) as well as in cigarette smokers. In contrast, elderly patients, patients with renal or hepatic dysfunction, and patients receiving  
5 concomitant drugs which inhibit the cytochrome P450 system should receive lower initial and maintenance doses, e.g., 75 mg. These dosages are simply guidelines and the actual dose selected for an individual patient will be determined by the attending physician based upon clinical conditions and using methods well-known in the art. Flupirtine may be provided in either a single or multiple dosage regimen, e.g., a patient may take 100 mg of flupirtine orally three times a day.

10

#### *Dosage Forms and Route of Administration*

Any route of administration and dosage form is compatible with the present invention and flupirtine may be administered as either the sole active agent or in combination with other therapeutically active drugs. Although compositions suitable for oral delivery are preferred, other  
15 routes that may be used include peroral, internal, pulmonary, rectal, nasal, vaginal, lingual, transdermal, intravenous, intraarterial, intramuscular, intraperitoneal, intracutaneous and subcutaneous routes. Specific dosage forms include tablets, pills, capsules, powders, aerosols, suppositories, skin patches, parenterals, and oral liquids including oil aqueous suspensions, solutions and emulsions. Sustained release dosage forms may also be used. All dosage forms  
20 may be prepared using methods that are standard in the art (see e.g., Remington's Pharmaceutical Sciences, 16th ed., A. Oslo Editor, Easton PA (1980)). Specific guidance for the preparation of dosage forms for various routes of delivery is provided by U.S. patents 4,668,684; 5,503,845; and 5,284,861.

25

Flupirtine may be used in conjunction with any of the vehicles and excipients commonly employed in pharmaceutical preparations, e.g., talc, gum arabic, lactose, starch, magnesium stearate, cocoa butter, aqueous or non-aqueous solvents, oils, paraffin derivatives, glycols, etc.

Coloring and flavoring agents may also be added to preparations, particularly to those for oral administration. Solutions can be prepared using water or physiological compatible organic  
30 solvents such as ethanol, 1,2-propylene glycol, polyglycols, dimethyl sulfoxide, fatty alcohols, triglycerides, partial esters of glycerine and the like. Parenteral compositions containing flupirtine

may be prepared using conventional techniques and include sterile isotonic saline, water, 1,3-butanediol, ethanol, 1,2-propylene glycol, polyglycols mixed with water, Ringer's solution, etc.

#### *Method of Treatment*

5 Generally, a patient diagnosed as having fibromyalgia may be initially given a relatively low dose of flupirtine, e.g., 50 mg per day, in order to determine whether any adverse side-effects are experienced. This is particularly important in cases where a patient is taking other medications or has clinical characteristics that suggest that they may not be able to tolerate high drug dosages. Although flupirtine is relatively safe when used at dosages lower than 600 mg per day, a number  
10 of side effects have been reported. Among these are dizziness, drowsiness, pruritus, dry mouth, and, less frequently, nausea, depression, sleep disturbance, and headache. If adverse effects are not experienced by the patient, dosage may be gradually increased until a satisfactory alleviation of the symptoms associated with fibromyalgia, particularly pain, is achieved. Since flupirtine is non-addictive, treatment may be safely maintained over a prolonged period of time.

15 The daily dose of flupirtine may be administered as a single tablet or capsule, but it is generally preferable to divide the daily dosage into two or more separate aliquots. Alternatively, a patient may simply take flupirtine as needed, up to the maximum tolerated daily dosage. Flupirtine administration may be combined with the administration of other therapeutically active  
20 agents, such as antidepressants, depending upon the individual needs of a patient.

#### *Treatment of Diseases or Conditions Related to Fibromyalgia*

In addition its use in the treatment of fibromyalgia, flupirtine may be used in exactly the manner described above for the treatment of a number of diseases and conditions related to  
25 fibromyalgia. Included among such diseases and conditions are fibrositis, chronic fatigue syndrome, myofascial pain syndrome, soft tissue rheumatism, non-articular rheumatism, idiopathic muscle pain syndrome, chronic widespread musculoskeletal pain, major depression associated with musculoskeletal pain, somatization disorder, somatoform pain disorder, Lyme disease with concurrent fibromyalgia, rheumatoid arthritis with concurrent fibromyalgia,  
30 generalized tendomyopathy, eosinophilia-myalgia syndrome, and stiff-man syndrome.

*Advantages of Flupirtine Over Opiates*

Patients receiving flupirtine do not acquire a tolerance to its analgesic effects the way that patients receiving opiates do. Thus, there is no need for a continual increase in dosage once an optimum has been found. Flupirtine is non-addictive and the abrupt discontinuation of its administration does not produce withdrawal symptoms. In addition, flupirtine is less sedating than opiate drugs and, unlike opiates, it exhibits muscle relaxant properties and is not associated with cardiovascular side effects.

Despite these differences in adverse effects, flupirtine has an analgesic action in patients with fibromyalgia and related conditions that is roughly comparable to the effect produced by opiates, and, in some instances, flupirtine may be superior. Thus, flupirtine has a therapeutic utility comparable to, or better than, some of the most effective antinociceptive agents known without comparable risks and side effects.

All references cited herein are fully incorporated by reference. Having now fully described the invention, it will be understood by those of skill in the art that the invention may be performed within a wide and equivalent range of conditions, parameters and the like, without affecting the spirit or scope of the invention or embodiment thereof.

***What is Claimed is:***

1. A method of treating a patient for fibromyalgia which comprises:

5        administering an amount of flupirtine to said patient effective to reduce or eliminate one or more symptoms associated with said fibromyalgia.

2. The method of claim 1, wherein said flupirtine is administered orally at a dose of between 50 mg and 900 mg per day, calculated on the basis of the free-base form of flupirtine.

10

3. The method of claim 2, wherein said flupirtine is administered at a dose of between 100 mg and 400 mg per day, calculated on the basis of the free-base form of flupirtine.

4. The method of claim 1, wherein said flupirtine is administered parenterally at a dose of  
15        between 50 mg and 900 mg per day, calculated on the basis of the free-base form of flupirtine.

5. The method of claim 4, wherein said flupirtine is administered at a dose of between 100 mg and 400 mg per day, calculated on the basis of the free-base form of flupirtine.

20        6. The method of claim 1, wherein said flupirtine is administered transdermally at a dosage of between 50 mg and 900 mg per day, calculated on the basis of the free-base form of flupirtine.

7. The method of claim 6, wherein said flupirtine is administered at a dosage of between 100 mg and 400 mg per day, calculated on the basis of the free-base form of flupirtine.

25

8. A method of treating a patient for a disease or condition related to fibromyalgia by administering flupirtine to said patient at a dosage of between 50 mg and 900 mg of flupirtine per day, calculated on the basis of the free-base form of flupirtine, and wherein said disease or condition is selected from the group consisting of: fibrositis; chronic fatigue syndrome;  
30        myofascial pain syndrome; soft tissue rheumatism; non-articular rheumatism; idiopathic muscle pain syndrome; chronic widespread musculoskeletal pain; major depression associated with

musculoskeletal pain; somatization disorder; somatoform pain disorder; Lyme disease with concurrent fibromyalgia; rheumatoid arthritis with concurrent fibromyalgia; generalized tendomyopathy; eosinophilia-myalgia syndrome; and stiff-man syndrome.

5 9. The method of claim 8, wherein said disease or condition is fibrositis.

10. The method of claim 8, wherein said disease or condition is chronic fatigue syndrome.

11. The method of claim 8, wherein said disease or condition is myofascial pain syndrome.

10

12. The method of claim 8, wherein said disease or condition is soft tissue rheumatism.

13. The method of claim 8, wherein said disease or condition is non-articular rheumatism.

15 14. The method of claim 8, wherein said disease or condition is idiopathic muscle pain syndrome.

15. The method of claim 8, wherein said disease or condition is chronic widespread musculoskeletal pain.

20

16. The method of claim 8, wherein said disease or condition is major depression associated with musculoskeletal pain.

17. The method of claim 8, wherein said disease or condition is somatization disorder.

25

18. The method of claim 8, wherein said disease or condition is somatoform pain disorder.

19. The method of claim 8, wherein said disease or condition is Lyme disease with concurrent fibromyalgia.

30

20. The method of claim 8, wherein said disease or condition is rheumatoid arthritis with concurrent fibromyalgia.
21. The method of claim 8, wherein said disease or condition is generalized tendomyopathy.
- 5 22. The method of claim 8, wherein said disease or condition is eosinophilia-myalgia syndrome.
23. The method of claim 8, wherein said disease or condition is stiff-man syndrome.